

BIMS 8601 Homework #1: Due in class Thursday, February 24, 2022

Stefan Bekiranov

February 18, 2022

1. Problem 1 (25 points):

- (a) Show that the Poisson distribution can be derived as the limit of the binomial distribution as the number of trials, n , approaches infinity and the probability of success on each trial, p approaches zero in such a way that $np = \lambda$. (Hint: For $n \rightarrow \infty$ use Sterling's approximation $n! \sim n^n e^{-n} \sqrt{2\pi n}$ and $(1 - \frac{\lambda}{n})^n \rightarrow e^{-\lambda}$.)
- (b) Show that the normal distribution can be derived as the limit of the binomial distribution as the number of trials, n , approaches infinity and the probability of success on each trial, p is not close to 0 or 1. Steps of derivation: (1) apply Sterling's approximation (2) collect terms and take the natural log of both sides of the equation (3) use a change of variables $\delta = k - np$ (4) apply the expansion $\ln(1 + x) \sim x - \frac{1}{2}x^2$ (5) assume δ is much smaller than n keeping lowest order terms and (6) apply the exponential function to both sides of the equation.

2. Problem 2 (25 points):

Let $Y = \sum_i^n X_i$ where the X_i are independent Bernoulli random variables that take the value 1 and 0 with probability p and $1 - p$, respectively. Calculate the expectation and variance of Y : $E(Y)$ and $Var(Y)$. How do you expect Y to be distributed? Briefly explain your answer and why this makes sense.

3. Problem 3 (25 points):

Calculate the expectation and variance of a random variable assumed to follow the (a) Poisson and (b) normal distribution. Feel free to use the moment generating function for the Poisson distribution $M(t) = e^{\lambda(e^t - 1)}$ and the normal distribution $M(t) = e^{\mu t + \sigma^2 t^2 / 2} = e^{\mu t + \sigma^2 t^2 / 2}$.

4. Problem 4 (25 points):

Prove the Central Limit Theorem by showing that the moment generating function of $Z_n = S_n / \sigma \sqrt{n}$ (where $S_n = \sum_{i=1}^n X_i$ and the X_i are independent random variables with mean 0 and variance σ^2 and moment generating function $M(t)$) tends to the moment generating function of the standard normal distribution as $n \rightarrow \infty$. Execute the following steps:

- (a) Calculate the moment generating function of Z_n , $M_{Z_n}(t)$, in terms of the moment generating function of the X_i , $M(t)$. Use the rules of how to calculate moment generating functions of sums of random variables and linear functions of random variables shown in class.
- (b) Use the fact that $M(s)$ has a Taylor series expansion $M(s) = M(0) + sM'(0) + \frac{1}{2}s^2M''(0) + \dots$ and the definition of expectation and variance in terms of $M'(0)$ and $M''(0)$, respectively.
- (c) Use the fact that as $n \rightarrow \infty$, $(1 + \frac{\lambda}{n})^n \rightarrow e^\lambda$.
- (d) Arrive at the final result, $M_{Z_n}(t) \rightarrow e^{t^2/2}$ as $n \rightarrow \infty$.